



A bioaugmentation approach to pesticide bioremediation

pesticide degradation and crop protection

Vlaiculescu, Alexandru; Jensen, Birgit; Varrone, Cristiano

DOI (link to publication from Publisher):
[10.13140/RG.2.2.15175.44964](https://doi.org/10.13140/RG.2.2.15175.44964)

Creative Commons License
Unspecified

Publication date:
2019

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Vlaiculescu, A., Jensen, B., & Varrone, C. (2019). *A bioaugmentation approach to pesticide bioremediation: pesticide degradation and crop protection*. Poster presented at Symposium on Bioengineering in Food and Feed Production. 13. Nov. 2019. DTU Bioengineering. Kg. Lyngby. Denmark, Lyngby, Denmark.
<https://doi.org/10.13140/RG.2.2.15175.44964>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/337284313>

A BIOAUGMENTATION APPROACH TO PESTICIDE BIOREMEDIATION: PESTICIDE DEGRADATION AND CROP PROTECTION

Poster · November 2019

DOI: 10.13140/RG.2.2.15175.44964

CITATIONS

0

READS

144

3 authors:



Alexandru Vlaiculescu

Aalborg University

2 PUBLICATIONS 0 CITATIONS

SEE PROFILE



Birgit Jensen

University of Copenhagen

56 PUBLICATIONS 1,189 CITATIONS

SEE PROFILE



Cristiano Varrone

Aalborg University

170 PUBLICATIONS 885 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



VegQURE [View project](#)



Linking biohydrogen and biodiesel production (EU FP7 STF China project) [View project](#)

A BIOAUGMENTATION APPROACH TO PESTICIDE BIOREMEDIATION: PESTICIDE DEGRADATION AND CROP PROTECTION

Alexandru Vlaiculescu¹, Birgit Jensen², Cristiano Varrone¹

¹Aalborg University Copenhagen, Department of Chemistry and Bioscience, Section of Sustainable Biotechnology

²University of Copenhagen; Department of Plant and Environmental Sciences; Section for Microbial Ecology and Biotechnology

BACKGROUND

- A pesticide is defined as any substance or mixture of substances intended for preventing, destroying, repelling, or mitigating pests ^[1].
- The main function of pesticides is crop protection. It is estimated that diseases, insects and weeds destroy between 31-42% of all crops produced worldwide, and the annual worldwide crop loss from plant diseases is about 220 billion dollars ^[2].
- Many organisms can degrade pesticides. Eukaryotic organisms, like many plants, animals, or fungi, use this ability to neutralize their harmful effects, while bacteria typically break down the pesticide to assimilate its nutrients. Bacteria are accounted for most of the pesticide biodegradation. Their ability to transfer genes horizontally, combined with a fast evolution rate, enables them to quickly adapt and extend the biodegradation potential of a new mutant to a whole bacterial community ^[3].



15 Most Abundant Genus	Sludge	Step 1	Step 2	Step 3	Step 4	Step 5	Step 6	% of Read Abundance
<i>Pseudomonas</i>	0.00	63.98	55.97	53.96	54.97	42.57	0.33	
<i>Hyphomicrobiaceae</i>	0.00	0.01	0.20	1.14	11.93	17.51	31.06	
<i>Methyloceanibacter</i>	50.84	0.15	0.10	0.14	0.01	0.06	0.00	
<i>Streptococcus</i>	0.00	9.83	13.67	9.61	0.78	0.52	0.00	
<i>Xanthobacteraceae</i>	0.00	0.00	0.01	0.06	0.23	1.33	31.61	
<i>Rhizobiaceae</i>	0.00	0.53	1.30	3.06	10.65	4.66	4.72	
<i>Shinella</i>	0.00	1.90	4.93	6.61	7.37	1.77	0.10	
<i>Parapedobacter</i>	0.00	2.17	3.02	3.59	1.39	9.73	0.66	
<i>Vagococcus</i>	0.00	8.06	5.30	2.92	0.14	0.85	0.01	
<i>Sphingomonadaceae</i>	0.00	1.33	2.15	4.57	4.00	1.49	0.93	
<i>Lactococcus</i>	0.00	3.37	2.81	1.75	0.24	1.25	0.00	
<i>Gammaproteobacteria</i>	8.30	0.00	0.01	0.02	0.00	0.00	0.00	
<i>Brevundimonas</i>	0.00	2.42	2.65	2.24	0.47	0.22	0.03	
<i>Devosiaceae</i>	0.00	1.13	2.02	2.19	0.91	0.31	0.96	
<i>Rhodospseudomonas</i>	0.00	0.00	0.00	0.00	0.01	0.27	6.31	

List of the 15 most abundant genus in the enrichment, and their abundance (%). The enrichment strategy lasted for 11 weeks, following 3 initial 1 week steps, a 4th 2 weeks step and two weeks 5th and 6th step.

It can be observed that a major change occurred between step 5 and step 6 of the enrichment, with the variability decreasing, and some species that were predominant until the fifth step (ex. *Pseudomonas*) almost disappear in the 6th step. Aside from the decrease in *Pseudomonas*, there is a high increase in the *Hyphomicrobiaceae* and *Xanthobacteraceae* genera.

Since the 5th step and the 6th step of enrichment are very different in variability, both will be utilized in the experiments.


PRELIMINARY RESULTS

Bacterial Degradation of Dicamba, Dimethoate and Malathion




Testing of 6 bacterial isolates in M9 media supplemented with Dicamba 50 mg/L, working volume 20 mL.

Dicamba Preliminary Results



Calibration curve of the HPLC method for the pesticide Dicamba.

The first step of determining the accuracy of the method was performed, obtaining an R square value of 0.9999. Since Dicamba is a herbicide, and it is highly soluble in water, it was chosen as the first pesticide to test.



Degradation of 60 mg/L Dicamba in M9 media. The results have been normalized and calculated through comparison with the control samples. The nomenclature of the organisms is: HS- Hungarian Soil, DIC- isolates from the enrichment performed with Dicamba, DIM- isolates from the enrichment performed with Dimethoate, IDEO- *Ideonella* sp., SLU- Sludge samples, MAL- isolates from the enrichment performed with Malathion. In this case, most organisms used were isolated from the Dicamba enrichment.

Dicamba Preliminary Conclusions

Compared to the control sample, one organism degraded 3.87 mg/L of Dicamba, HSDIC24, an isolate that was enriched on Dicamba, while most others seem to have not degraded the herbicide. This represents only the first batch of tested organisms, approximately 60 more isolates shall be tested in order to determine if they can degrade the herbicide Dicamba.

RESEARCH QUESTIONS

- Is it possible to find efficient pesticide degrading organisms in contaminated soils and sludge?
- Do pesticide-degrading organisms also exhibit crop protection function?
- Can pesticide-degrading microorganisms improve plant growth and prevent toxicity of pesticides present?



METHODS

- Enrichment** of environmental samples in M9 media supplemented with pesticides (Dicamba, Dimethoate, Malathion).

- Isolation** of strains at the end of the enrichment.

- Pesticide Degradation Assay:** Degradation efficiency of the previously isolated strains is assessed through a 3 week experiment. The organisms are incubated at 30 degrees C in M9 media supplemented with Dicamba/Dimethoate/Malathion. Determination of the degraded amount is performed through HPLC analysis.

- Development of a **Defined Microbial Consortium (DMC)**. The DMC will be formed from the best degraders found from testing the isolated strains.

- The Crop Protection Function:** the effect of pesticide-degrading bacteria on plant growth parameters and their ability to control fungal plant pathogens will be tested in plant assays.

